

Derivative Operators in Metric and Geometric Structures

V. V. Fernández¹, A. M. Moya², W. A. Rodrigues Jr.¹

¹Institute of Mathematics, Statistics and Scientific Computation

IMECC-UNICAMP CP 6065

13083-970 Campinas, SP, Brazil

e-mail: walrod@ime.unicamp.br ; virginvelfe@accessplus.com.ar

²Department of Mathematics, University of Antofagasta, Antofagasta, Chile

e-mail: mmoya@uantof.cl

February 1, 2008

Abstract

This paper (the seventh paper in a series of eight) continues the development of our theory of multivector and extensor calculus on smooth manifolds. Here we deal first with the concepts of ordinary Hodge coderivatives, duality identities, and Hodge coderivative identities. Then, we recall the concept of a Levi-Civita geometric structure and the concepts of Levi-Civita and gauge derivatives. New formulas that are important in the Lagrangian theory of multivector and extensor fields are obtained. We introduce also the concept of *covariant* Hodge coderivative. We detail how all these concepts are related.

Contents

1	Introduction	2
2	Ordinary Hodge Coderivatives	2
2.1	Duality Identities	5
2.2	Hodge Duality Identities	6

2.3	Ordinary Hodge Coderivative Operators	6
3	Levi-Civita Geometric Structure	7
3.1	Levi-Civita Derivatives	9
3.2	Gauge Derivatives	11
4	Covariant Hodge Coderivative	14
5	Conclusions	16

1 Introduction

This is the seventh paper in a series of eight. Here, we continue the development of our theory of multivector and extensor calculus on smooth manifolds. We introduce in Section 2 the ordinary Hodge coderivatives, duality identities, and Hodge coderivative identities. In Section 3 we recall the concept of a Levi-Civita geometric structure and the concepts of Levi-Civita and *gauge* derivatives. Several important formulas that appear in the Lagrangian formulation of the theory of multivector and extensor fields on smooth manifolds are obtained.¹ In Section 4 we introduce the new concept of *covariant* Hodge coderivative. We study in details how all these important concepts are related. Finally in Section 5 we present our conclusions.

2 Ordinary Hodge Coderivatives

Let U be an open subset of U_o , and let (U, g) be a *metric structure* on U . Let us take any pair of *reciprocal frame fields* on U , say $(\{e_\mu\}, \{e^\mu\})$, i.e., $e_\mu \cdot e^\nu = \delta_\mu^\nu$. In particular, the *fiducial frame field*, namely $\{b_\mu\}$, due to its *orthonormality*, i.e., $b_\mu \cdot b_\nu = \delta_{\mu\nu}$, is a *self-reciprocal frame field*, i.e., $b^\mu = b_\mu$. We note also that $\{b_\mu\}$ is an *ordinarily constant frame field* on U , i.e.,

$$a \cdot \partial_o b_\mu = 0, \text{ for each } \mu = 1, \dots, n. \quad (1)$$

¹In [1] we gave a preliminary presentation of the Lagrangian theory of multivector and extensor fields on Minkowski spacetime. A more general theory based on the developments of the present series of papers is in preparation.

Associated to $(\{e_\mu\}, \{e^\mu\})$, the smooth pseudoscalar field on U , namely τ , defined by

$$\tau = \sqrt{e_\wedge \cdot e_\wedge} e^\wedge, \quad (2)$$

where $e_\wedge = e_1 \wedge \dots \wedge e_n \in \mathcal{M}^n(U)$ and $e^\wedge = e^1 \wedge \dots \wedge e^n \in \mathcal{M}^n(U)$, is said to be the *standard volume pseudoscalar field* for the *local coordinate system* (U_o, ϕ_o) .

Such $\tau \in \mathcal{M}^n(U)$ has the fundamental property

$$\tau \cdot \tau = \tau \lrcorner \tilde{\tau} = \tau \tilde{\tau} = 1. \quad (3)$$

It follows from the obvious result $e_\wedge \cdot e^\wedge = 1$.

>From Eq.(3) we can get an expansion formula for smooth pseudoscalar fields on U , i.e.,

$$I = (I \cdot \tau) \tau. \quad (4)$$

In particular, because of the obvious properties $b_\wedge \cdot b_\wedge = 1$ and $b^\wedge = b_\wedge$, the standard volume pseudoscalar field associated to $\{b_\mu\}$ is just $b_\wedge$. It will be called the *canonical volume pseudoscalar field* for (U_o, ϕ_o) .

We emphasize that $b_\wedge$ is an *ordinarily constant* smooth pseudoscalar field on U , i.e.,

$$a \cdot \partial_o b_\wedge = 0. \quad (5)$$

Eq.(5) can be proved by using Eq.(1) and the Leibnitz rule for the exterior product of smooth multivector fields.

By using Eq.(5) and the general Leibnitz rule for $a \cdot \partial_o$ we can deduce the remarkable property

$$a \cdot \partial_o (b_\wedge * X) = b_\wedge * (a \cdot \partial_o X), \quad (6)$$

where $*$ means either exterior product or any canonical product of smooth multivector fields.

On the other hand we have that all $(\{e_\mu\}, \{e^\mu\})$ must be necessarily an *extensor-deformation* of $\{b_\mu\}$. This statement means that there exists a non-singular smooth $(1, 1)$ -extensor field on U , say ε , such that

$$e_\mu = \varepsilon(b_\mu), \quad (7)$$

$$e^\mu = \varepsilon^*(b_\mu), \text{ for each } \mu = 1, \dots, n. \quad (8)$$

Then, by putting Eq.(7) and Eq.(8) into Eq.(2), we have that

$$\tau = (\underline{\varepsilon}(b_\wedge) \cdot \underline{\varepsilon}(b_\wedge))^{1/2} \underline{\varepsilon}^*(b_\wedge) = (\det^2 [\varepsilon] b_\wedge \cdot b_\wedge)^{1/2} \det^{-1} [\varepsilon] b_\wedge = \text{sgn}(\det[\varepsilon]) b_\wedge,$$

i.e.,

$$\tau = \pm b_{\wedge}. \quad (9)$$

>From Eq.(5) and Eq.(6), by taking into account Eq.(9), we have two remarkable properties

$$a \cdot \partial_o \tau = 0, \quad (10)$$

$$a \cdot \partial_o (\tau * X) = \tau * (a \cdot \partial_o X), \text{ for all } X \in \mathcal{M}(U). \quad (11)$$

Associated to $(\{e_\mu\}, \{e^\mu\})$, the smooth pseudoscalar field on U , namely τ_g , defined by

$$\tau_g = \sqrt{\left| e_{\wedge} \cdot_g e_{\wedge} \right|} e^{\wedge} = \sqrt{|\det[g]|} \tau, \quad (12)$$

will be said to be a *metric volume pseudoscalar field* for (U_o, ϕ_o) .

Such $\tau_g \in \mathcal{M}^n(U)$ satisfies the basic property

$$\tau_g \cdot_{g^{-1}g} \tau_g = \tau_g \lrcorner_g \tilde{\tau}_g = \tau_{gg^{-1}g} \tilde{\tau}_g = (-1)^q. \quad (13)$$

In order to prove it we should recall that $\text{sgn}(\det[g]) = (-1)^q$, where q is the *number of negative eigenvalues* of g .

An expansion formula for smooth pseudoscalar fields on U can be also obtained from Eq.(13), i.e.,

$$I = (-1)^q (I \cdot_{g^{-1}g} \tau_g) \tau_g. \quad (14)$$

Associated to τ , the smooth extensor field on U , namely $\star$, defined by

$$\star X = \tilde{X} \lrcorner \tau = \tilde{X} \tau, \quad (15)$$

will be called the *standard Hodge extensor field* on U .

Such $\star$ is *non-singular* and its inverse $\star^{-1}$ is given by

$$\star^{-1} X = \tau \lrcorner \tilde{X} = \tau \tilde{X}. \quad (16)$$

By using a property analogous to that result given by Eq.(11) we can easily prove that the standard Hodge extensor field is *ordinarily constant*, i.e.,

$$a \cdot \partial_o \star = 0. \quad (17)$$

Also,

$$a \cdot \partial_o \star^{-1} = 0. \quad (18)$$

Associated to τ_g , the smooth extensor field on U , namely $\star_g$, defined by

$$\star_g X = \tilde{X} \lrcorner_g \tau = \tilde{X} \lrcorner_{g^{-1}g} \tau = \sqrt{|\det[g]|} \underline{g}^{-1}(\tilde{X}) \lrcorner \tau = \sqrt{|\det[g]|} \underline{g}^{-1}(\tilde{X}) \tau, \quad (19)$$

will be called the *metric Hodge extensor field* on U . Of course, it is associated to the metric structure (U, g) .

Such $\star_g$ is also non-singular and its inverse, namely $\star_g^{-1}$, is given by

$$\begin{aligned} \star_g^{-1} X &= (-1)^q \tau \lrcorner_{gg^{-1}} \tilde{X} = (-1)^q \tau \lrcorner_{gg^{-1}} \tilde{X}, \\ &= (-1)^q \sqrt{|\det[g]|} \tau \lrcorner \underline{g}^{-1}(\tilde{X}) = (-1)^q \sqrt{|\det[g]|} \tau \underline{g}^{-1}(\tilde{X}). \end{aligned} \quad (20)$$

2.1 Duality Identities

We present in this subsection two interesting and useful formulas which relate the *ordinary curl* $\partial_o \wedge$ to the *ordinary contracted divergence* $\partial_o \lrcorner$.

i. For all $X \in \mathcal{M}(U)$ it holds

$$\tau(\partial_o \wedge X) = (-1)^{n+1} \partial_o \lrcorner (\tau X). \quad (21)$$

Proof

We will use the so-called duality identity $I(a \wedge Y) = (-1)^{n+1} a \lrcorner (IY)$, where $a \in \mathcal{U}_o$, $I \in \bigwedge^n \mathcal{U}_o$ and $Y \in \bigwedge \mathcal{U}_o$. By recalling the known identities $\partial_a \wedge (a \cdot \partial_o X) = \partial_o \wedge X$ and $\partial_a \lrcorner (a \cdot \partial_o X) = \partial_o \lrcorner X$, and using Eq.(11), we get

$$\begin{aligned} \tau(\partial_o \wedge X) &= \tau(\partial_a \wedge (a \cdot \partial_o X)) = (-1)^{n+1} \partial_a \lrcorner (\tau(a \cdot \partial_o X)), \\ &= (-1)^{n+1} \partial_a \lrcorner (a \cdot \partial_o (\tau X)) = (-1)^{n+1} \partial_o \lrcorner (\tau X). \blacksquare \end{aligned}$$

ii. For all $X \in \mathcal{M}(U)$ it holds

$$\tau_{g^{-1}}(\partial_o \wedge X) = \frac{(-1)^{n+1}}{\det[g]} \underline{g}(\partial_o \lrcorner (\tau X)). \quad (22)$$

Proof

It can be deduced by using the identity² $I_{g^{-1}} Y = \det^{-1}[g] \underline{g}(IY)$ and Eq.(21). $\blacksquare$

²In order to prove it we should use the following identities: $I_{g^{-1}} Y = I \underline{g}^{-1}(Y)$, $X \lrcorner \underline{t}^{-1}(Y) = \underline{t}^\dagger(\underline{t}^*(X) \lrcorner Y)$ and $\underline{t}^{-1}(I) = \det^{-1}[\underline{t}] I$, and so forth.

2.2 Hodge Duality Identities

Now, we present two noticeable identities which relate the curl $\partial_o \wedge$ to the contracted divergence $\partial_o \lrcorner$ involving the standard and the metric Hodge extensor fields $\star$ and $\star_g$.

i. For all $X \in \mathcal{M}(U)$ it holds

$$\star^{-1}(\partial_o \wedge (\star X)) = -\partial_o \lrcorner \widehat{X}. \quad (23)$$

Proof

We will use the duality identity given by Eq.(21). By using Eq.(16) and Eq.(15), and recalling the identities $\widetilde{\partial_o \wedge Y} = \partial_o \wedge \overline{Y}$ and $\overline{XY} = \overline{Y} \overline{X}$, and the obvious property $\tau \overline{\tau} = (-1)^n$, we get

$$\begin{aligned} \star^{-1}(\partial_o \wedge (\star X)) &= \tau(\widetilde{\partial_o \wedge (\tilde{X} \tau)}) = \tau(\partial_o \wedge (\overline{\tau} \widehat{X})), \\ &= (-1)^{n+1} \partial_o \lrcorner (\tau \overline{\tau} \widehat{X}) = -\partial_o \lrcorner \widehat{X}. \blacksquare \end{aligned}$$

ii. For all $X \in \mathcal{M}(U)$ it holds

$$\star_g^{-1}(\partial_o \wedge (\star_g X)) = -\frac{1}{\sqrt{|\det[g]|}} \underline{g}(\partial_o \lrcorner (\sqrt{|\det[g]|} \underline{g}^{-1}(\widehat{X}))). \quad (24)$$

Proof

We will use the duality identity given by Eq.(22). A straightforward calculation using Eq.(20) and Eq.(19) allows us to get

$$\begin{aligned} \star_g^{-1}(\partial_o \wedge (\star_g X)) &= (-1)^q \sqrt{|\det[g]|} \tau_{g^{-1}}(\widetilde{\partial_o \wedge (\tilde{X}_{g^{-1}g} \tau)}) \\ &= (-1)^q \sqrt{|\det[g]|} \tau_{g^{-1}} \partial_o \wedge (\sqrt{|\det[g]|} \overline{\tau} \underline{g}^{-1}(\widehat{X})) \\ &= (-1)^q \sqrt{|\det[g]|} \frac{(-1)^{n+1}}{\det[g]} \underline{g}(\partial_o \lrcorner (\sqrt{|\det[g]|} \tau \overline{\tau} \underline{g}^{-1}(\widehat{X}))), \\ &= -\frac{1}{\sqrt{|\det[g]|}} \underline{g}(\partial_o \lrcorner (\sqrt{|\det[g]|} \underline{g}^{-1}(\widehat{X}))). \blacksquare \end{aligned}$$

2.3 Ordinary Hodge Coderivative Operators

We introduce the so-called *standard Hodge derivative operator* $\delta : \mathcal{M}(U) \rightarrow \mathcal{M}(U)$ such that

$$\delta X = \star^{-1}(\partial_o \wedge (\star \widehat{X})). \quad (25)$$

Its basic property is

$$\delta X = -\partial_o \lrcorner X. \quad (26)$$

It immediately follows from the Hodge duality identity given by Eq.(23).

We introduce the so-called *metric Hodge coderivative operator* $\delta_g : \mathcal{M}(U) \rightarrow \mathcal{M}(U)$ such that

$$\delta_g X = \star_g^{-1} (\partial_o \wedge (\star_g \widehat{X})). \quad (27)$$

It satisfies the basic property

$$\delta_g X = -\frac{1}{\sqrt{|\det[g]|}} g(\partial_o \lrcorner (\sqrt{|\det[g]|} \underline{g}^{-1}(X))), \quad (28)$$

which is an immediate consequence of the Hodge duality identity given by Eq.(24).

3 Levi-Civita Geometric Structure

We recall [2] that the *Levi-Civita connection field* is the smooth vector elementary 2-extensor field on U , namely λ , defined by

$$\lambda(a, b) = \frac{1}{2} g^{-1} \circ (a \cdot \partial_o g)(b) + \omega_0(a) \times_g b, \quad (29)$$

where ω_0 is the smooth $(1, 2)$ -extensor field on U given by

$$\omega_0(a) = -\frac{1}{4} \underline{g}^{-1} (\partial_b \wedge \partial_c) a \cdot ((b \cdot \partial_o g)(c) - (c \cdot \partial_o g)(b)). \quad (30)$$

Such ω_0 satisfies

$$\omega_0(a) \times_g b \cdot c = \frac{1}{2} a \cdot ((b \cdot \partial_o g)(c) - (c \cdot \partial_o g)(b)). \quad (31)$$

The *open set* U endowed with λ and g , namely (U, λ, g) , is a *geometric structure* on U , a statement that means that *Levi-Civita parallelism structure* (U, λ) is compatible with the metric structure (U, g) . Or equivalently, the pair of *a-DCDO's* associated to (U, λ) , namely (D_a^+, D_a^-) , is g -compatible.

The *Levi-Civita a-DCDO's* D_a^+ and D_a^- are defined by

$$D_a^+ X = a \cdot \partial_o X + \Lambda_a(X), \quad (32)$$

$$D_a^- X = a \cdot \partial_o X - \Lambda_a^\dagger(X). \quad (33)$$

Note that Λ_a is the so-called *generalized* of λ_a . The latter is the so-called *a-directional connection field* associated to λ , given by $\lambda_a(b) = \lambda(a, b)$.

We present now two pairs of noticeable properties of λ .

i. The *scalar divergence* of $\lambda_a(b)$ with respect to the first variable, namely $\partial_a \cdot \lambda_a(b)$, and the *curl* of $\lambda_a^\dagger(b)$ with respect to the first variable, namely $\partial_a \wedge \lambda_a^\dagger(b)$, are given by

$$\partial_a \cdot \lambda_a(b) = \frac{1}{\sqrt{|\det[g]|}} b \cdot \partial_o \sqrt{|\det[g]|}, \quad (34)$$

$$\partial_a \wedge \lambda_a^\dagger(b) = 0. \quad (35)$$

Proof

In order to prove the first result we will use the formula $\tau^*(\partial_n) \cdot (a \cdot \partial_o \tau)(n) = \det^{-1}[\tau] a \cdot \partial_o \det[\tau]$, valid for all non-singular smooth $(1, 1)$ -extensor field τ . By using the symmetry property $(g^{-1})^\dagger = g^{-1}$ and Eq.(31), we can write

$$\begin{aligned} \partial_a \cdot \lambda_a(b) &= \frac{1}{2} \partial_a \cdot (g^{-1} \circ (a \cdot \partial_o g)(b) + \partial_a \cdot (\omega_0(a) \times_g b)) \\ &= \frac{1}{2} g^{-1}(\partial_a) \cdot (a \cdot \partial_o g)(b) + \omega_0(g^{-1}(\partial_a)) \times_g b \cdot_g a \\ &= \frac{1}{2} g^{-1}(\partial_a) \cdot (b \cdot \partial_o g)(a) = \frac{1}{2} \frac{1}{\det[g]} b \cdot \partial_o \det[g]. \end{aligned}$$

Then, recalling the identity $(f)^{-1} b \cdot \partial_o f = 2(|f|)^{-1/2} b \cdot \partial_o (|f|)^{1/2}$ valid for all non-zero $f \in \mathcal{S}(U)$, the expected result immediately follows.

To prove the second result we only need to take into account the *symmetry property* $\lambda_a(b) = \lambda_b(a)$. We have

$$\partial_a \wedge \lambda_a^\dagger(b) = \partial_a \wedge \partial_n(n \cdot \lambda_a^\dagger(b)) = \partial_a \wedge \partial_n(\lambda_a(n) \cdot b) = 0. \blacksquare$$

ii. The *left contracted divergence* of $\Lambda_a(X)$ with respect to a , namely $\partial_a \lrcorner \Lambda_a(X)$, and the *curl* of $\Lambda_a^\dagger(X)$ with respect to a , namely $\partial_a \wedge \Lambda_a^\dagger(X)$, are given by

$$\partial_a \lrcorner \Lambda_a(X) = \frac{1}{\sqrt{|\det[g]|}} (\partial_o \sqrt{|\det[g]|}) \lrcorner X, \quad (36)$$

$$\partial_a \wedge \Lambda_a^\dagger(X) = 0. \quad (37)$$

Proof

In order to prove the first ones we will use the multivector identities $v \lrcorner (X \wedge Y) = (v \lrcorner X) \wedge Y + \widehat{X} \wedge (v \lrcorner Y)$ and $X \lrcorner (Y \lrcorner Z) = (X \wedge Y) \lrcorner Z$, where $v \in \mathcal{U}_o$ and $X, Y, Z \in \bigwedge \mathcal{U}_o$. A straightforward calculation using Eq.(34) allows us to get

$$\begin{aligned} \partial_a \lrcorner \Lambda_a(X) &= \partial_a \cdot \lambda_a(\partial_n)(n \lrcorner X) - \lambda_a(\partial_n) \wedge (\partial_a \lrcorner (n \lrcorner X)) \\ &= \frac{1}{\sqrt{|\det[g]|}} (\partial_n \cdot \partial_o \sqrt{|\det[g]|})(n \lrcorner X) - \lambda_a(\partial_n) \wedge ((\partial_a \wedge n) \lrcorner X), \\ &= \frac{1}{\sqrt{|\det[g]|}} \partial_n (n \cdot \partial_o \sqrt{|\det[g]|}) \lrcorner X - \lambda_{\partial_a}(\partial_n) \wedge ((a \wedge n) \lrcorner X). \end{aligned}$$

Then, recalling the identity $\partial_n(n \cdot \partial_o Y) = \partial_o Y$ and the symmetry property $\lambda_a(b) = \lambda_b(a)$, we get the required result.

The proof of the second property follows immediately by using Eq.(35), and recalling that the adjoint of generalized is equal to the generalized of adjoint, i.e., $\Lambda_a^\dagger(X) = \lambda_a^\dagger(\partial_n) \wedge (n \lrcorner X)$. ■

3.1 Levi-Civita Derivatives

We introduce now the *canonical covariant divergence operator*, $D^+ \lrcorner : \mathcal{M}(U) \rightarrow \mathcal{M}(U)$ such that

$$D^+ \lrcorner X = \partial_a \lrcorner (D_a^+ X), \quad (38)$$

i.e., $D^+ \lrcorner X = e^\mu \lrcorner (D_{e_\mu}^+ X) = e_\mu \lrcorner (D_{e^\mu}^+ X)$, where $(\{e_\mu\}, \{e^\mu\})$ is any pair of *canonical reciprocal frame fields* on U .

Its basic property is

$$D^+ \lrcorner X = \frac{1}{\sqrt{|\det[g]|}} \partial_o \lrcorner (\sqrt{|\det[g]|} X). \quad (39)$$

Indeed, using the identity $\partial_a \lrcorner (a \cdot \partial_o X) = \partial_o \lrcorner X$ and Eq.(36) into the definition given by Eq.(32), we get

$$\begin{aligned} D^+ \lrcorner X &= \partial_a \lrcorner (a \cdot \partial_o X) + \partial_a \lrcorner \Lambda_a(X), \\ &= \partial_o \lrcorner X + \frac{1}{\sqrt{|\det[g]|}} (\partial_o \sqrt{|\det[g]|}) \lrcorner X. \end{aligned}$$

So, by recalling the identity $\partial_o \lrcorner (fY) = (\partial_o f) \lrcorner X + f(\partial_o \lrcorner X)$, for all $f \in \mathcal{S}(U)$ and $Y \in \mathcal{M}(U)$, we can get the proof for this remarkable property.

The so-called *metric covariant divergence*, *covariant curl* and *metric covariant gradient operators*, namely $D^- \lrcorner_{g^{-1}}$, $D^- \wedge$ and $D^-_{g^{-1}}$, map also smooth multivector fields to smooth multivector fields and are defined by

$$D^- \lrcorner_{g^{-1}} X = \partial_a \lrcorner_{g^{-1}} (D_a^- X) = g^{-1}(\partial_a) \lrcorner (D_a^- X), \quad (40)$$

$$D^- \wedge X = \partial_a \wedge (D_a^- X), \quad (41)$$

$$D^-_{g^{-1}} X = \partial_a_{g^{-1}} (D_a^- X). \quad (42)$$

The relationship among these operators is given by

$$D^-_{g^{-1}} X = D^- \lrcorner_{g^{-1}} X + D^- \wedge X. \quad (43)$$

The basic properties of the metric covariant divergence are

$$D^- \lrcorner_{g^{-1}} X = \underline{g}(D^+ \lrcorner \underline{g}^{-1}(X)), \quad (44)$$

$$D^- \lrcorner_{g^{-1}} X = \frac{1}{\sqrt{|\det[g]|}} \underline{g}(\partial_o \lrcorner (\sqrt{|\det[g]|} \underline{g}^{-1}(X))). \quad (45)$$

Eq.(44) follows from the fundamental property $D_a^- \underline{g}(X) = \underline{g}(D_a^+ X)$ which holds for any g -compatible pair of a -DCDO's, by using the identity $X \lrcorner \underline{t}(Y) = \underline{t}(\underline{t}^\dagger(X) \lrcorner Y)$. Eq.(45) is deduced by using Eq.(44) and Eq.(39).

A remarkable property which follows from Eq.(45) is

$$D^- \lrcorner_{g^{-1}} (D^- \lrcorner_{g^{-1}} X) = 0. \quad (46)$$

In order to prove it we should use the known identity $\partial_o \lrcorner (\partial_o \lrcorner Y) = 0$.

By comparing Eq.(28) and Eq.(45) we get

$$D^- \lrcorner_{g^{-1}} X = -\delta_g X. \quad (47)$$

The basic property for the covariant curl is

$$D^- \wedge X = \partial_o \wedge X. \quad (48)$$

It immediately follows by using the identity $\partial_a \wedge (a \cdot \partial_o X) = \partial_o \wedge X$ and Eq.(37) into the definition given by Eq.(33).

The four *derivative-like operators* defined by Eq.(38), Eq.(40), Eq.(41) and Eq.(42) will be called the *Levi-Civita derivatives*. They are involved in three useful identities which are used in the *Lagrangian theory of multivector fields*. These are,

$$\begin{aligned} & (\partial_o \wedge X) \cdot_{g^{-1}} Y + X \cdot_{g^{-1}} (D^- \lrcorner_{g^{-1}} Y) \\ &= \frac{1}{\sqrt{|\det[g]|}} \partial_o \cdot (\sqrt{|\det[g]|} \partial_n (n \wedge X) \cdot_{g^{-1}} Y), \end{aligned} \quad (49)$$

$$\begin{aligned} & (D^- \lrcorner_{g^{-1}} X) \cdot_{g^{-1}} Y + X \cdot_{g^{-1}} (\partial_o \wedge Y) \\ &= \frac{1}{\sqrt{|\det[g]|}} \partial_o \cdot (\sqrt{|\det[g]|} \partial_n (n \lrcorner_{g^{-1}} X) \cdot_{g^{-1}} Y), \end{aligned} \quad (50)$$

$$\begin{aligned} & (D^- \lrcorner_{g^{-1}} X) \cdot_{g^{-1}} Y + X \cdot_{g^{-1}} (D^- \lrcorner_{g^{-1}} Y) \\ &= \frac{1}{\sqrt{|\det[g]|}} \partial_o \cdot (\sqrt{|\det[g]|} \partial_n (n \lrcorner_{g^{-1}} X) \cdot_{g^{-1}} Y). \end{aligned} \quad (51)$$

3.2 Gauge Derivatives

Let h be a gauge metric field for g [3]. This statement means that there is a smooth $(1, 1)$ -extensor field h such that $g = h^\dagger \circ \eta \circ h$, where η is an orthogonal metric field with the same signature as g . As we know, associated to a Levi-Civita a -DCDO's, namely (D_a^+, D_a^-) , there must be an unique η -compatible pair of a -DCDO's, namely $(\mathbb{D}_a^+, \mathbb{D}_a^-)$, given by the following formulas

$$\mathbb{D}_{ha}^+ X = \underline{h}(D_a^+ \underline{h}^{-1}(X)), \quad (52)$$

$$\mathbb{D}_{h^*a}^- X = \underline{h}^*(D_a^- \underline{h}^\dagger(X)). \quad (53)$$

These equations say that $(\mathbb{D}_{ha}^+, \mathbb{D}_{h^*a}^-)$ is the h -deformation of (D_a^+, D_a^-) . So, $\mathbb{D}_{ha}^+$ and $\mathbb{D}_{h^*a}^-$ will be called the *gauge covariant derivatives* associated to D_a^+ and D_a^- .

We present here two noticeable properties of $\mathbb{D}_{ha}^+$.

i. For all $a, b, c \in \mathcal{V}(U)$, it holds

$$(\mathbb{D}_{ha}^+ b) \cdot_\eta c = [h(a), h^{-1}(b), h^{-1}(c)]. \quad (54)$$

Proof

By using Eq.(52), the identity $(D_a^+ b) \cdot c = \left\{ \begin{smallmatrix} c \\ a, b \end{smallmatrix} \right\}$, and the definition of the Christoffel operator of second kind, i.e., $\left\{ \begin{smallmatrix} c \\ a, b \end{smallmatrix} \right\} = [a, b, g^{-1}(c)]$, we can write

$$(\mathbb{D}_{ha}^+ b) \cdot_\eta c = D_a^+(h^{-1}(b)) \cdot h^\dagger \circ \eta(c) = \left\{ \begin{smallmatrix} h^\dagger \circ \eta(c) \\ h(a), h^{-1}(b) \end{smallmatrix} \right\} = [h(a), h^{-1}(b), g^{-1} \circ h^\dagger \circ \eta(c)],$$

hence, by recalling that $g^{-1} = h^{-1} \circ \eta \circ h^*$, the required result immediately follows. ■

ii. There exists a smooth $(1, 2)$ -extensor field on U_o , namely Ω_0 , such that

$$\mathbb{D}_a^+ X = a \cdot \partial_o X + \Omega_0(a) \times_\eta X. \quad (55)$$

Such Ω_0 is given by

$$\Omega_0(a) = -\frac{1}{2} \underline{\eta}(\partial_b \wedge \partial_c)[a, h^{-1}(b), h^{-1}(c)]. \quad (56)$$

Proof

Firstly, we must prove a particular case of the above property, i.e.,

$$\mathbb{D}_a^+ b = a \cdot \partial_o b + \Omega_0(a) \times_\eta b. \quad (57)$$

We will use the following properties of the Christoffel operator of first kind $[a, b, c] - [a, c, b] = 2[a, b, c] - a \cdot \partial_o(b \cdot_g c)$, and $[a, b + b', c] = [a, b, c] + [a, b', c]$ and $[a, fb, c] = f[a, b, c] + (a \cdot \partial_o f)b \cdot_g c$. A straightforward calculation allows

us to get

$$\begin{aligned}
\Omega_0(a) \times_{\eta} b \cdot_{\eta} c &= (b \wedge c) \cdot_{\eta} \Omega_0(a) = \frac{1}{2} (b \wedge c) \cdot (\partial_p \wedge \partial_q) [a, h^{-1}(p), h^{-1}(q)] \\
&= \frac{1}{2} \det \begin{bmatrix} b \cdot \partial_p & b \cdot \partial_q \\ c \cdot \partial_p & c \cdot \partial_q \end{bmatrix} [\dots] \\
&= \frac{1}{2} b \cdot \partial_p c \cdot \partial_q [\dots] - b \cdot \partial_q c \cdot \partial_p [\dots] \\
&= \frac{1}{2} b \cdot \partial_p c \cdot \partial_q ([a, h^{-1}(p), h^{-1}(q)] - [a, h^{-1}(q), h^{-1}(p)]) \\
&= \frac{1}{2} b \cdot \partial_p c \cdot \partial_q (2[\dots] - a \cdot \partial_o (h^{-1}(p) \cdot_g h^{-1}(q))) \\
&= b \cdot \partial_p c \cdot \partial_q (p \cdot b_{\mu} [a, h^{-1}(b_{\mu}), h^{-1}(c)] + (a \cdot \partial_o p) \cdot_{\eta} c) \\
&= b \cdot b_{\mu} [a, h^{-1}(b_{\mu}), h^{-1}(c)] + (a \cdot \partial_o b) \cdot_{\eta} c - (a \cdot \partial_o b) \cdot_{\eta} c \\
&= [a, h^{-1}(b), h^{-1}(c)] - (a \cdot \partial_o b) \cdot_{\eta} c,
\end{aligned}$$

hence, by using Eq.(54), the particular case given by Eq.(57) immediately follows. Now, we can indeed prove the general case of the above property .

As we can see, the a -directional connection field for $(\mathbb{D}_a^+, \mathbb{D}_a^-)$ is given by $b \mapsto \Omega_0(a) \times_{\eta} b$. Then, its generalized (extensor field) must be given by $X \mapsto (\Omega_0(a) \times_{\eta} \partial_b) \wedge (b \lrcorner X)$. But, by recalling the noticeable identity $(B \times_{\eta} \partial_b) \wedge (b \lrcorner X) = B \times_{\eta} X$, where $B \in \bigwedge^2 \mathcal{U}_o$ and $X \in \bigwedge \mathcal{U}_o$, we find that it can be written as $X \mapsto \Omega_0(a) \times_{\eta} X$. ■

We introduce now the *gauge covariant divergence*, *gauge covariant curl* and *gauge covariant gradient operators*, namely $\mathbb{D}_{\eta}^{-} \lrcorner$, $\mathbb{D}_{\eta}^{-} \wedge$ and $\mathbb{D}_{\eta}^{-}$. They all map smooth multivector fields to smooth multivector fields and are defined by

$$\mathbb{D}_{\eta}^{-} \lrcorner X = h^*(\partial_a) \lrcorner_{\eta} (\mathbb{D}_{h^*a}^{-} X), \quad (58)$$

$$\mathbb{D}_{\eta}^{-} \wedge X = h^*(\partial_a) \wedge (\mathbb{D}_{h^*a}^{-} X), \quad (59)$$

$$\mathbb{D}_{\eta}^{-} X = h^*(\partial_a) (\mathbb{D}_{h^*a}^{-} X). \quad (60)$$

It is obvious that the relationship among them is given by

$$\mathbb{D}_{\eta}^{-} X = \mathbb{D}_{\eta}^{-} \lrcorner X + \mathbb{D}_{\eta}^{-} \wedge X. \quad (61)$$

Their basic properties are given by the following *golden formulas*

$$\underline{h}^*(D^- \underset{g^{-1}}{\lrcorner} X) = \mathbb{D}^- \underset{\eta}{\lrcorner} \underline{h}^*(X), \quad (62)$$

$$\underline{h}^*(D^- \wedge X) = \mathbb{D}^- \wedge \underline{h}^*(X), \quad (63)$$

$$\underline{h}^*(D^- \underset{g^{-1}}{\lrcorner} X) = \mathbb{D}^- \underset{\eta}{\lrcorner} \underline{h}^*(X). \quad (64)$$

These formulas can be proved by using the golden formula deduced in [3], $\underline{h}^*(X \underset{g^{-1}}{*} Y) = \underline{h}^*(X) \underset{\eta}{*} \underline{h}^*(Y)$, where $X, Y \in \bigwedge \mathcal{U}_o$, and $\underset{g^{-1}}{*}$ means either exterior product or any g^{-1} -product of smooth multivector fields, and analogously for $\underset{\eta}{*}$. It is also necessary to take into account the master formulas $g = h^\dagger \circ \eta \circ h$ and $g^{-1} = h^{-1} \circ \eta \circ h^*$, and the relationship between $\mathbb{D}_{ha}^+$ and $\mathbb{D}_{h^*a}^-$, i.e., $\underline{\eta}(\mathbb{D}_{ha}^+ \underline{\eta}(X)) = \mathbb{D}_{h^*a}^- X$.

>From Eq.(45) and Eq.(48) by using Eq.(62) and Eq.(63) we find the interesting identities

$$\mathbb{D}^- \underset{\eta}{\lrcorner} X = \frac{1}{\det[h]} \underline{\eta} \circ h(\partial_o \lrcorner (\det[h] \underline{h}^{-1} \circ \underline{\eta}(X))), \quad (65)$$

$$\mathbb{D}^- \wedge X = \underline{h}^*(\partial_o \wedge \underline{h}^\dagger(X)). \quad (66)$$

4 Covariant Hodge Coderivative

Let (U, γ, g) be a *geometric structure* on U , and let us denote by $(\mathcal{D}_a^+, \mathcal{D}_a^-)$ the *g-compatible* pair of *a-DCDO's* associated to (U, γ, g) . We will present two noticeable properties which are satisfied by the second *a-DCDO*.

- i. τ is a *covariantly constant* smooth pseudoscalar field on U , i.e.,

$$\mathcal{D}_a^- \tau = 0. \quad (67)$$

Proof

We will use the basic property and the expansion formula given by Eq.(13) and Eq.(14). By using the Ricci-like theorem for $\mathcal{D}_a^-$, we have that

$$(\mathcal{D}_a^- \tau) \underset{g^{-1}}{\cdot} \underset{g}{\tau} + \tau \underset{g}{\cdot} \underset{g^{-1}}{(\mathcal{D}_a^- \tau)} = 0,$$

i.e., $(\mathcal{D}_a^- \tau) \underset{g^{-1}}{\cdot} \underset{g}{\tau} = 0$. Then, $\mathcal{D}_a^- \tau = (-1)^q ((\mathcal{D}_a^- \tau) \underset{g^{-1}}{\cdot} \underset{g}{\tau}) \underset{g}{\tau} = 0$. ■

ii. For all $X \in \mathcal{M}(U)$ it holds

$$\mathcal{D}_a^-(\tau_{g g^{-1}} * X) = \tau_{g g^{-1}} * \mathcal{D}_a^-(X), \quad (68)$$

where $*_{g^{-1}}$ means either exterior product or any g^{-1} -product of smooth multivector fields.

Proof

It can be deduced by using Eq.(67) and the general Leibnitz rule for $\mathcal{D}_a^-$. ■

We introduce the so-called *metric covariant divergence*, *covariant curl* and *metric covariant gradient operators*, namely $\mathcal{D}^- \lrcorner_{g^{-1}}$, $\mathcal{D}^- \wedge$ and $\mathcal{D}^-_{g^{-1}}$, which map smooth multivector fields to smooth multivector fields. They are defined by

$$\mathcal{D}^- \lrcorner_{g^{-1}} X = \partial_a \lrcorner_{g^{-1}} (\mathcal{D}_a^- X), \quad (69)$$

$$\mathcal{D}^- \wedge X = \partial_a \wedge (\mathcal{D}_a^- X), \quad (70)$$

$$\mathcal{D}^-_{g^{-1}} X = \partial_a \lrcorner_{g^{-1}} (\mathcal{D}_a^- X). \quad (71)$$

The relationship among them is given by

$$\mathcal{D}^-_{g^{-1}} X = \mathcal{D}^- \lrcorner_{g^{-1}} X + \mathcal{D}^- \wedge X. \quad (72)$$

Now, we will present two noticeable duality identities between $\mathcal{D}^- \wedge$ and $\mathcal{D}^- \lrcorner_{g^{-1}}$. One of them involves the metric Hodge extensor field $\star_g$.

iii. For all $X \in \mathcal{M}(U)$ it holds

$$\tau_{g g^{-1}} (\mathcal{D}^- \wedge X) = (-1)^{n+1} \mathcal{D}^- \lrcorner_{g^{-1}} (\tau_{g g^{-1}} X). \quad (73)$$

Proof

We will use the g^{-1} -duality identity $I_{g^{-1}}(a \wedge Y) = (-1)^{n+1} a \lrcorner_{g^{-1}} (I_{g^{-1}} Y)$, where $a \in \mathcal{U}_o$, $I \in \bigwedge^n \mathcal{U}_o$ and $Y \in \bigwedge \mathcal{U}_o$. A straightforward calculation by taking into account Eq.(68) gives

$$\begin{aligned} \tau_{g g^{-1}} (\mathcal{D}^- \wedge X) &= (-1)^{n+1} \partial_a \lrcorner_{g^{-1}} (\tau_{g g^{-1}} (\mathcal{D}_a^- X)) \\ &= (-1)^{n+1} \partial_a \lrcorner_{g^{-1}} (\mathcal{D}_a^- (\tau_{g g^{-1}} X)), \\ &= (-1)^{n+1} \mathcal{D}^- \lrcorner_{g^{-1}} (\tau_{g g^{-1}} X). \blacksquare \end{aligned}$$

iv. For all $X \in \mathcal{M}(U)$ it holds

$$\star_g^{-1}(\mathcal{D}^- \wedge (\star_g X)) = -\mathcal{D}^- \lrcorner_{g^{-1}} \hat{X}. \quad (74)$$

Proof

We will use the duality identity given by Eq.(73). By using Eq.(20) Eq.(19), and recalling the identities $\widetilde{\mathcal{D}^- \wedge X} = \mathcal{D}^- \wedge \overline{X}$ and $\overline{XY} = \overline{Y} \overline{X}$, and the obvious property $\tau_{gg^{-1}g} \overline{\tau} = (-1)^{n+q}$, we get

$$\begin{aligned} \star_g^{-1}(\mathcal{D}^- \wedge (\star_g X)) &= (-1)^q \tau_{gg^{-1}}(\widetilde{\mathcal{D}^- \wedge (\widetilde{\widetilde{X}}_{g^{-1}g} \tau)}) \\ &= (-1)^q \tau_{gg^{-1}}(\mathcal{D}^- \wedge (\overline{\tau}_{gg^{-1}} \hat{X})) \\ &= (-1)^q (-1)^{n+1} \mathcal{D}^- \lrcorner_{g^{-1}} (\tau_{gg^{-1}g} \overline{\tau}_{gg^{-1}} \hat{X}), \\ &= -\mathcal{D}^- \lrcorner_{g^{-1}} \hat{X}. \blacksquare \end{aligned}$$

We introduce now the *covariant Hodge coderivative operator*, namely Δ_g , as defined by $\Delta_g : \mathcal{M}(U) \rightarrow \mathcal{M}(U)$ such that

$$\Delta_g X = \star_g^{-1}(\mathcal{D}^- \wedge (\star_g \hat{X})). \quad (75)$$

Its basic property is

$$\mathcal{D}^- \lrcorner_{g^{-1}} X = -\Delta_g X, \quad (76)$$

which follows trivially from Eq.(74).

5 Conclusions

In this paper (the fourth paper in a series of five) we continued the development of our theory of multivector and extensor calculus on smooth manifolds. We introduced first the concepts of ordinary Hodge coderivatives, duality identities, and Hodge coderivative identities. Then, we recalled the concept of a Levi-Civita geometric structure and the concepts of Levi-Civita and gauge derivatives. We also introduced the new concept of *covariant* Hodge coderivative. We studied in details how all these important concepts

are related and obtained several formulas which play a key role in the Lagrangian theory of multivector and extensor fields on arbitrary manifolds. We believe that our results clarify many misconceptions in earlier literature on the subject.

Acknowledgments: V. V. Fernández and A. M. Moya are very grateful to Mrs. Rosa I. Fernández who gave to them material and spiritual support at the starting time of their research work. This paper could not have been written without her inestimable help.

References

- [1] Fernández, V. V., Moya, A. M., and Rodrigues, W. A., Jr., Lagrangian Formalism for Multiform Fields on Minkowski Spacetime, *Int. J. Theor. Phys.* **40**, 299-313 (2001).
- [2] Rodrigues, W. A., Jr., Fernández, V. V., Moya, A. M., and Metric Compatible Covariant Derivatives, submitted for publication.
- [3] Moya, A. M., Fernández, V. V., and Rodrigues, W. A., Jr., Metric and Gauge Extensors, submitted for publication.